



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A61K 39/395, 39/00, A01N 63/00, 1/02, 1/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/27433</b> <b>(43) International Publication Date:</b> 18 May 2000 (18.05.00)
<b>(21) International Application Number:</b> PCT/US99/24012 <b>(22) International Filing Date:</b> 9 November 1999 (09.11.99)  <b>(30) Priority Data:</b> 60/107,657                      9 November 1998 (09.11.98)                      US  <b>(71) Applicant (for all designated States except US):</b> IDEC PHARMACEUTICALS CORPORATION [US/US]; 11011 Torreyana Road, San Diego, CA 92121 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> GRILLO-LOPEZ, Antonio, J. [US/US]; P.O. Box 3797, Rancho Santa Fe, CA 92067 (US). LEONARD, John, E. [US/US]; 3440 Camino Alegre, Carlsbad, CA 92009 (US).  <b>(74) Agents:</b> GESS, E., Joseph et al.; Burns, Doane, Swecker & Mathis L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> CHIMERIC ANTI-CD20 ANTIBODY TREATMENT OF PATIENTS RECEIVING BMT OR PBSC TRANSPLANTS		
<b>(57) Abstract</b> <p>The use of a chimeric anti-CD20 antibody, RITUXAN®, as an <i>in vivo</i> or <i>in vitro</i> purging agent, of patients receiving bone marrow or peripheral blood stem cell transplant during treatment of B-cell-related malignancies, e.g., B-cell lymphomas or leukemias, is disclosed. Such purging may enhance engraftment and/or prevent disease relapse in such patients.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## CHIMERIC ANTI-CD20 ANTIBODY TREATMENT OF PATIENTS RECEIVING BMT OR PBSC TRANSPLANTS

5

**FIELD OF THE INVENTION**

The use of an anti-CD20 antibody or a fragment thereof as an *in vitro* or *in vivo* purging agent in patients receiving bone marrow transplant or peripheral blood stem cell transplant is disclosed.

10

**BACKGROUND OF THE INVENTION**

The use of antibodies to the CD20 antigen as diagnostic and/or therapeutic agents for B-cell lymphoma has previously been reported. CD20 is a useful marker or target for B-cell lymphomas as this antigen is expressed at very high densities on the surface of malignant B-cells, i.e., B-cells wherein unabated proliferation can lead to B-cell lymphomas.

15

CD20 or Bp35 is a B-lymphocyte-restricted differentiation antigen that is expressed during early pre-B-cell development and remains until plasma cell differentiation. It is believed by some that the CD20 molecule may regulate a step in the B-cell activation process which is required for cell cycle initiation and differentiation. Moreover, as noted, CD20 is usually expressed at very high levels on neoplastic ("tumor") B-cells.

20

Previous reported therapies involving anti-CD20 antibodies have involved the administration of a therapeutic anti-CD20 antibody either alone or in conjunction with a second radiolabeled anti-CD20 antibody, or a chemotherapeutic agent.

In fact, the Food and Drug Administration has approved the therapeutic use of one such therapeutic anti-CD20 antibody, RITUXAN®, for use in relapsed and previously treated low-grade non-Hodgkin's lymphoma (NHL).

Also, the use of RITUXAN® in combination with a radiolabeled murine anti-  
5 CD20 antibody has been suggested for the treatment of B-cell lymphoma.

However, while anti-CD20 antibodies and, in particular, RITUXAN®, have been reported to be effective for treatment of B-cell lymphomas, such as non-Hodgkin's lymphoma, the treated patients are often subject to disease relapse. Therefore, it would be beneficial if more effective antibody treatments could be  
10 developed. More specifically, it would be advantageous if other therapeutic applications of anti-CD20 antibodies were discovered. Also, it would be helpful if current treatment protocols for B-cell lymphoma were improved, which prevented or further reduced disease relapse.

## 15 **BRIEF DESCRIPTION OF THE INVENTION**

Thus, it is an object of the invention to improve the problems of prior treatments of B-cell-related diseases, e.g., B-Cell lymphomas and leukemias, in particular the problem of disease relapse after disease treatment.

More specifically, it is an object of the invention to reduce the incidence of  
20 disease relapse in patients with B-cell-related diseases receiving bone marrow or peripheral blood stem cell transplants by the use of an anti-CD20 antibody as an *in vitro* and/or *in vivo* purging agent prior, concurrent, and/or after transplant.

It is an even more specific object of the invention to use RITUXAN® as an  
*in vitro* and/or *in vivo* purging agent prior, concurrent and/or after bone marrow or  
25 peripheral blood stem cell transplant.

### DETAILED DESCRIPTION OF THE INVENTION

A significant problem associated with the treatment of diseases involving B-cells and other cells that express CD20 antigen, including B-cell lymphomas and leukemias, is the problem of disease relapse after treatment.

5           The exact cause for disease relapse is unclear. However, it is known that such relapse may occur even in patients that receive aggressive therapeutic intervention, e.g., high dosages of chemotherapeutic agents, cytokines, radiation, and/or antibody. While the exact cause of relapse remains unclear, it is speculated by some researchers that disease relapse may occur because the patient may still harbor low numbers of  
10       diseased cells even after aggressive therapy. Also, it is speculated that bone marrow transplant or peripheral blood stem cell transplanted tissue may itself be contaminated by diseased cells that express the CD20 antigen, e.g., B-cell lymphoma cells. Therefore, transplant of such tissues may unwittingly introduce diseased cells, and thereby actually increase the risk of disease relapse.

15           As discussed, the present invention seeks to prevent or reduce the incidence of disease in patients receiving transplanted bone marrow or peripheral blood stem cells by treating the transplanted bone marrow or peripheral blood stem cells with an amount of an anti-CD20 antibody or fragment thereof effective to purge the transplanted tissue of disease-causing CD20 antigen-expressing cells. Such purging  
20       may be effected *in vitro* and/or *in vivo*.

For example, bone marrow or peripheral blood stem cells may be contacted in tissue culture with an anti-CD20 antibody prior to transplant. In the preferred embodiment such antibody will comprise a chimeric, primate, primatized®, humanized or human anti-CD20 antibody, preferably RITUXAN®.

25           Alternatively, or in conjunction with such *in vitro* purging, the patient may be treated concurrent or subsequent to bone marrow or peripheral blood stem cell

transplant with an amount of an anti-CD20 antibody effective to purge (*in vivo*) or at least reduce the number of disease causing cells that express CD20 antigen that may be present in the transplant.

Similarly, the antibody used for *in vivo* purging will preferably comprise a  
5 chimeric, humanized, primate, primatized®, or human anti-CD20 antibody,  
preferably RITUXAN®. This *in vivo* purging may be effected simultaneous or  
substantially contemporaneous to bone marrow or peripheral blood stem cell  
transplant. Preferably, such purging will be effected within a week or more,  
preferably within 1 to 12 hours after transplant. However, such purging can be  
10 effected up to about 1 to 100 days after transplant. In the preferred embodiment, *in*  
*vivo* purging will be effected within about 1 month after transplant, more preferably  
within about one week after transplant, and most preferably within about 1 to 12  
hours after transplant.

As noted above, the subject *in vivo* or *in vitro* purging of CD20 antigen-  
15 expressing cells will desirably be effected in patients that have previously been  
treated in an effort to eradicate disease causing B-cells, or other CD20 antigen-  
expressing cells involved in disease. Such treatment methods include, by way of  
example, cytokine therapy, antibody therapies (e.g., RITUXAN® or other antibodies  
targeted to B-cells), chemotherapy and/or radiation therapy, e.g., whole body  
20 irradiation, radioimmunotherapy.

In an especially preferred embodiment, the subject *in vitro* or *in vivo* purging  
will be effected in patients that have previously been treated with RITUXAN®  
and/or radioimmunotherapy that receive an autologous bone marrow or peripheral  
blood stem cell transplant after RIT and/or RITUXAN® therapy.

For example, patients that have a B-cell-related disease, e.g., a B-cell lymphoma or leukemia, will have their bone marrow or peripheral blood stem cells collected prior to therapeutic treatment. This will be effected by known methods.

The patient will then be subjected to an aggressive therapeutic regimen, e.g.,  
5 administration of RITUXAN®, or a radiolabeled antibody that is specific to an antigen expressed by the tumor cells, whole body irradiation, and/or a chemotherapeutic or cytokine. This therapeutic regimen will be effected under conditions that are hypothetically designed to eradicate any B-cell or other CD20 antigen-expressing tumor cells that may be present.

10 After such treatment has been completed, the bone marrow or peripheral blood stem cells, which optionally may be treated *in vitro* with an anti-CD20 antibody, e.g., RITUXAN®, to deplete CD20 expressing cells, is then transplanted into the patient in order to reconstitute the immune system thereof.

Concurrently or shortly thereafter, the patient will be administered an amount  
15 of an anti-CD20 antibody, e.g., RITUXAN®, effective to purge any disease causing cells that may be present in the bone marrow or peripheral blood stem cell transplant. An effective dosage will typically comprise from about 0.01 to about 3.0 mg/kg body weight. A preferred dosage will comprise from about .1 to about 20 mg/kg, more preferably from about .1 to about 5.0 mg/kg, administered within about one week of  
20 transplant.

The subject *in vitro* and/or *in vivo* purging will reduce the risk of relapse in many B-cell-related diseases, e.g., B-cell lymphomas and leukemias such as non-Hodgkin's lymphomas, chronic lymphocytic leukemia, etc., after treatment has been completed in patients receiving transplanted cells that potentially may be  
25 contaminated with disease-causing cells.

Also, the subject method should be well tolerated based on the relative non-toxicity of anti-CD20 antibodies, such as RITUXAN®, and therefore should not adversely impact engraftment of the transplanted autologous cells. In fact, it may act to promote engraftment of such transplant.

5       As noted in the preferred embodiment, the purging agent will comprise RITUXAN®. However, other anti-CD20 antibodies may be used, e.g., other chimeric, primate, primatized®, humanized or human antibodies. Also, antibody fragments may be used, e.g., Fv's, FAB, F(ab)', F(ab<sub>2</sub>)<sup>1</sup>, and aggregates thereof. In addition, antibodies and antibody fragments directed to other B cell surface  
10       markers, e.g., CD19, may also be used.

Methods for producing chimeric, primate, primatized®, humanized and human antibodies are well known in the art. See, e.g., U.S. Patent 5,530,101, issued to Queen et al, U.S. Patent 5,225,539, issued to Winter et al, U.S. Patents 4,816,397 and 4,816,567, issued to Boss et al, and Cabilly et al, respectively, all of which are  
15       incorporated by reference in their entirety.

The selection of human constant regions may be significant to the therapeutic efficacy of the subject anti-CD20 antibody. In the preferred embodiment, the subject anti-CD20 antibody will comprise human, gamma 1, or gamma 3 constant regions and, more preferably, human gamma 1 constant regions. The use of gamma 1 anti-  
20       CD20 antibodies as therapeutics is disclosed in U.S. Patent 5,500,362, issued to Robinson et al.

Methods for making human antibodies are also known and include, by way of example, production in SCID mice, and *in vitro* immunization.

As noted, a particularly preferred chimeric anti-CD20 antibody is  
25       RITUXAN®, which is a chimeric gamma 1 anti-human CD20 antibody. The complete amino acid and corresponding nucleic acid sequence for this antibody may



be found in U.S. Patent 5,736,137, which is incorporated by reference in its entirety. This antibody, which is produced in a proprietary CHO cell expression system commercialized by IDEC Pharmaceuticals Corporation, is made by a CHO cell transfectoma which was deposited on November 4, 1992, under the provisions of the Budapest Treaty at the American Type Culture Collection, located at 12301 Parklawn Drive, Rockville, MD 20852. This cell line was determined to be viable and will be replaced should it become non-viable during the term of deposit. This cell line was made irrevocably available upon issuance of the 5,736,137 patent and is available without restriction from the ATCC. This cell line will also be available without restriction during the lifetime of any patent that may issue based on this application.

The subject anti-CD20 antibody, when used as a purging agent, will be administered by various routes of administration, typically parenteral. This is intended to include intravenous, intramuscular, subcutaneous, rectal, vaginal, and administration with intravenous infusion being preferred.

The anti-CD20 antibody will be formulated for therapeutic usage by standard methods, e.g., by addition of pharmaceutically acceptable buffers, e.g., sterile saline, sterile buffered water, propylene glycol, and combinations thereof.

#### EXAMPLE

A single-arm pivotal study of Rituximab® infused at 375 mg/m<sup>2</sup> weekly times four was conducted in 166 patients with relapsed or refractory, low-grade or follicular NHL (International Working Formulation [IWF] Types A - D and REAL classification, small lymphocytic lymphoma, Follicular center, follicular Grades I, II, III). (McLaughlin P, Grillo-López A, Link B, Levy R, Czuczman M, Williams M, Heyman M, Bence-Bruckler I, White C, Cabanillas F, Jain V, Ho A, Lister J, Wey K, Shen D, Dallaire B. Rituximab® chimeric anti-CD20 monoclonal antibody therapy for relapsed indolent lymphoma: half of patients respond to a 4-dose

treatment program. Journal of Clinical Oncology 1998; 16:2825-2833.) Patients with tumor masses > 10 cm or with > 5000 lymphocytes/ $\mu$ L in the peripheral blood were excluded from this study. The median age was 58 years (105 men and 61 women) and the median number of prior treatments was three. Bone marrow involvement was present in 56% of 149 patients evaluated. Forty-five percent had  $\geq 2$  extranodal sites and 41% had bulky disease ( $\geq 5$  cm).

Complete response required the regression of all lymph nodes to  $< 1 \times 1$  cm<sup>2</sup> demonstrated on two occasions at least 28 days apart on neck, chest, abdomen, and pelvic CT scans, resolution of all symptoms and signs of lymphoma, and normalization of bone marrow, liver, and spleen. Partial response required a  $\geq 50\%$  decrease in the sum of the products of perpendicular measurements of lesions without any evidence of progressive disease for at least 28 days. Patients who did not achieve a CR or PR were considered non-responders, even if a net decrease ( $> 50\%$ ) of measurable disease was observed. Time to progression was measured from the first infusion until progression.

The overall response rate (ORR) was 48% with a 6% CR and a 42% PR rate (McLaughlin P, Grillo-López A, Link B, Levy R, Czuczman M, Williams M, Heyman M, Bence-Bruckler I, White C, Cabanillas F, Jain V, Ho A, Lister J, Wey K, Shen D, Dallaire B. Rituximab® chimeric anti-CD20 monoclonal antibody therapy for relapsed indolent lymphoma: half of patients respond to a 4-dose treatment program. Journal of Clinical Oncology 1998; 16:2825-2833). The median time to progression (TTP) for responders was 13.2 months and the median duration of response (DR) was 11.6 months. Twenty-two of 80 (28%) responders remain in ongoing remission at 20.9+ to 32.9+ months (McLaughlin P, Grillo-López A, Maloney D, Link B, Levy R, Czuczman M, Cabanillas F, Dallaire B, White C.

Efficacy controls in long-term follow-up of patients treated with rituximab for relapsed or refractory, low-grade or follicular NHL. *Blood* 1998; 92:414a-415a).

Administration of Rituximab® resulted in a rapid and sustained depletion of B-cells. Circulating B-cells were depleted within the first three doses with sustained  
5 depletion for up to six to nine months post-treatment in 83% of patients. Median B-cell levels returned to normal by 12 months following treatment. Although median NK cell counts remained unchanged, a positive correlation was observed between higher absolute NK cell counts at baseline and response to Rituximab® (Janakiraman N, McLaughlin P, White C, Maloney D, Shen D, Grillo-López A. Rituximab:  
10 Correlation between effector cells and clinical activity in NHL. *Blood* 1998; 92 (10 Suppl 1):337a).

Several baseline prognostic factors were analyzed to determine their correlation to response. Significantly, in 23 patients relapsed after ABMT or PBSC, the ORR was 78% versus 43% in patients who did not undergo prior high-dose  
15 therapy ( $p < 0.01$ ). This suggests that anti-CD20 treatment may effectively be used to purge CD20 antigen-expressing cells *in vivo* when administered following transplantation. Moreover, because patients who receive prior high dose therapy accompanied a bone marrow or peripheral stem cell transplantation appear to benefit more from subsequent Rituximab® therapy than those patients without prior therapy  
20 and transplantation, this suggests that a combined treatment protocol including bone marrow or stem cell transplantation provides a synergistic effect when compared to either single treatment alone.

Although the present invention has been described in some detail by way of illustration and example, for purposes of clarity and understanding it will be apparent  
25 that certain changes and modifications may be practical within the scope of the appended claims.

**WHAT IS CLAIMED IS:**

1. A method for reducing the risk of relapse of a B-cell-related disease in a patient receiving a bone marrow or peripheral blood stem cell transplant comprising treating said transplant *in vitro* and/or *in vivo* with an amount of an anti-CD20 antibody effective to reduce (purge) the number of disease-causing CD20 antigen-expressing cells therein.  
5
2. The method of Claim 1, wherein said disease is a B-cell lymphoma or leukemia.  
10
3. The method of Claim 1, wherein said purging is effected *in vivo* by administering RITUXAN® no later than about one month after transplant.
4. The method of Claim 1, wherein RITUXAN® is administered at a  
15 dosage ranging from about 0.1 to 20 mg/kg about one week after transplant.
5. The method of Claim 4, wherein the patient receiving the transplant has previously been treated under conditions designed to eradicate disease-causing B-cells.  
20
6. The method of Claim 4, wherein said patient has previously been subjected to a treatment protocol selected from the group consisting of whole body irradiation, RITUXAN® immunotherapy, chemotherapy, cytokine therapy, radioimmunotherapy, or a combination thereof.  
25

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/24012

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : A61K 39/395, 39/00; A01N 63/00, 1/02, 1/00

US CL : 424/93.7, 93.71, 133.1, 143.1, 144.1, 153.1, 155.1, 174.1, 435/2

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/93.7, 93.71, 133.1, 143.1, 144.1, 153.1, 155.1, 174.1, 435/2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MEDLINE, BIOSIS, EMBASE, WEST 1.2 search terms: author names, cd20, bone marrow, transplant, stem cells, pbse, c2b8, B cell, lymphoma

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,736,137 A (ANDERSON et al.) 07 April 1998, see entire document.	1-6



Further documents are listed in the continuation ... Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

10 JANUARY 2000

Date of mailing of the international search report

04 FEB 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RON SCHWADRON

Telephone No. (703) 308-0196